

PA DEPARTMENT OF HUMAN SERVICES
MAAC BRIEFING DOCUMENT
DUPIXENT (DUPILUMAB)

Proposed Effective Date: January 5, 2026

Revisions are noted with a ~~strikethrough~~ for deletions and **bold and underline** for additions.

I. Requirements for Prior Authorization of Dupixent (dupilumab)

A. Prescriptions That Require Prior Authorization

All prescriptions for Dupixent (dupilumab) must be prior authorized.

B. Revisions to Review of Documentation for Medical Necessity

In evaluating a request for prior authorization of a prescription for Dupixent (dupilumab), the determination of whether the requested prescription is medically necessary will take into account whether the beneficiary:

1. Is being treated for a diagnosis that is indicated in the U.S. Food and Drug Administration (FDA)-approved package labeling or a medically accepted indication; **AND**
2. Is age-appropriate according to FDA-approved package labeling, nationally recognized compendia, or peer-reviewed medical literature; **AND**
3. Is prescribed a dose that is consistent with FDA-approved package labeling, nationally recognized compendia, or peer-reviewed medical literature; **AND**
4. Is prescribed Dupixent (dupilumab) by or in consultation with an appropriate specialist (e.g., pulmonologist, allergist, immunologist, dermatologist, hematologist/oncologist, rheumatologist, etc.); **AND**
5. If currently using a different Monoclonal Antibody (MAB) – Anti-IL, Anti-IgE, Anti-TSLP, will discontinue the other MAB – Anti-IL, Anti-IgE, Anti-TSLP prior to starting Dupixent (dupilumab); **AND**
6. If currently using a different targeted systemic Immunomodulators, ~~Atopic Dermatitis~~ **Dermatologic** (e.g., Adbry [tralokinumab], Cibinqo [abrocitinib], **Nemluvio [nemolizumab]**, Rinvoq [upadacitinib]), will discontinue the other targeted systemic Immunomodulators, ~~Atopic Dermatitis~~ **Dermatologic** prior to starting Dupixent (dupilumab); **AND**
7. For a diagnosis of ~~moderate to severe~~ chronic atopic dermatitis, **both of the following:**
 - a. **Has atopic dermatitis associated with at least one of the following:**
 - i. **A body surface area of 10% or greater that is affected,**
 - ii. **Involvement of critical areas (e.g., face, feet, genitals, hands, intertriginous areas, scalp),**
 - iii. **Significant disability or impairment of physical, mental, or psychosocial**

PA DEPARTMENT OF HUMAN SERVICES
MAAC BRIEFING DOCUMENT
DUPIXENT (DUPILUMAB)

functioning

- b. Has a history of therapeutic failure of or a contraindication or an intolerance to **both** of the following:
 - i. **One** of the following:
 - a) For treatment of the face, skin folds, or other critical areas, a four-week trial of a low-potency topical corticosteroid
 - b) For treatment of other areas, a four-week trial of a medium-potency or higher topical corticosteroid
 - ii. An eight-week trial of a topical calcineurin inhibitor;

AND

- 8. For a diagnosis of asthma, **all** of the following:
 - a. Has asthma severity consistent with the FDA-approved indication for Dupixent (dupilumab) despite maximal therapeutic doses of or a contraindication or an intolerance to asthma controller drugs based on current national treatment guidelines for the diagnosis and management of asthma,
 - b. **One** of the following:
 - i. Has absolute blood eosinophil count ≥ 150 cells/microL
 - ii. Is dependent on oral corticosteroids,
 - c. Will use Dupixent (dupilumab) in addition to standard asthma controller drugs as recommended by current national treatment guidelines;

AND

- 9. For a diagnosis of eosinophilic esophagitis, has a history of therapeutic failure of or a contraindication or an intolerance to a proton pump inhibitor; **AND**
- 10. For a diagnosis of prurigo nodularis, **both** of the following:
 - a. Has a history of pruritis lasting at least six weeks
 - b. Has prurigo nodularis associated with at least **one** of the following:
 - i. ≥ 20 nodular lesions
 - ii. Significant disability or impairment of physical, mental, or psychosocial functioning;

AND

PA DEPARTMENT OF HUMAN SERVICES
MAAC BRIEFING DOCUMENT
DUPIXENT (DUPILUMAB)

11. **For a diagnosis of bullous pemphigoid, both of the following:**

a. **One of the following:**

- i. **Has a history of therapeutic failure of or a contraindication or an intolerance to systemic corticosteroids**
- ii. **Has corticosteroid-dependent disease**

b. **One of the following:**

- i. **Has a history of therapeutic failure of a corticosteroid-sparing therapy (e.g., doxycycline, dapson, methotrexate, mycophenolate, azathioprine)**
- ii. **Has a contraindication or an intolerance to corticosteroid-sparing therapies;**

AND

12. For all other diagnoses, has a history of therapeutic failure of or a contraindication or an intolerance to first line therapy(ies) if applicable according to consensus treatment guidelines; **AND**
13. If a prescription Dupixent (dupilumab) is for a quantity that exceeds the quantity limit, the determination of whether the prescription is medically necessary will also take into account the guidelines set forth in the Quantity Limits Chapter. See Quantity Limits List: <https://www.pa.gov/en/agencies/dhs/resources/pharmacy-services/quantity-limits-daily-dose-limits.html>.

NOTE: If the beneficiary does not meet the clinical review guidelines listed above but, in the professional judgment of the physician reviewer, the services are medically necessary to meet the medical needs of the beneficiary, the request for prior authorization will be approved.

FOR RENEWALS OF PRIOR AUTHORIZATION FOR DUPIXENT (DUPILUMAB): The determination of medical necessity of a request for renewal of a prior authorization for Dupixent (dupilumab) that was previously approved will take into account whether the beneficiary:

- 1. Is prescribed a dose that is consistent with FDA-approved package labeling, nationally recognized compendia, or peer-reviewed medical literature; **AND**
- 2. Is prescribed Dupixent (dupilumab) by or in consultation with an appropriate specialist (e.g., pulmonologist, allergist, immunologist, dermatologist, hematologist/oncologist, rheumatologist, etc.); **AND**
- 3. Has documented evidence of improvement in disease severity; **AND**

PA DEPARTMENT OF HUMAN SERVICES
MAAC BRIEFING DOCUMENT
DUPIXENT (DUPILUMAB)

4. For a diagnosis of asthma, **both** of the following:
- a. **One** of the following:
 - i. Has documented measurable evidence of improvement in the severity of the asthma condition
 - ii. Has reduction of oral corticosteroid dose while maintaining asthma control
 - b. Continues to use Dupixent (dupilumab) in addition to standard asthma controller drugs as recommended by current national treatment guidelines;

AND

5. If a prescription Dupixent (dupilumab) is for a quantity that exceeds the quantity limit, the determination of whether the prescription is medically necessary will also take into account the guidelines set forth in the Quantity Limits Chapter. See Quantity Limits List: <https://www.pa.gov/en/agencies/dhs/resources/pharmacy-services/quantity-limits-daily-dose-limits.html>.

NOTE: If the beneficiary does not meet the clinical review guidelines listed above but, in the professional judgment of the physician reviewer, the services are medically necessary to meet the medical needs of the beneficiary, the request for prior authorization will be approved.

C. Clinical Review Process

Prior authorization personnel will review the request for prior authorization and apply the clinical guidelines in Section B. above to assess the medical necessity of a prescription for Dupixent (dupilumab). If the guidelines in Section B. are met, the reviewer will prior authorize the prescription. If the guidelines are not met, the prior authorization request will be referred to a physician reviewer for a medical necessity determination. Such a request for prior authorization will be approved when, in the professional judgment of the physician reviewer, the services are medically necessary to meet the medical needs of the beneficiary.

D. References

1. Dupixent [package insert]. Tarrytown, NY: Regeneron Pharmaceuticals, Inc.: ~~October 2022~~ **June 2025**.
2. Howe W, Paller AS, Butala S. Treatment of atopic dermatitis (eczema). In: UpToDate [internet database]. Dellavalle RP, Levy ML, Fowler J, Corona R, eds. Waltham, MA: UpToDate. Updated July 25, 2023. Accessed August 1, 2023.
3. Lio PA. Management of severe atopic dermatitis (eczema) in children. In: UpToDate [internet database]. Dellavalle RP, Levy ML, Fowler J, Corona R, eds. Waltham, MA: UpToDate. Updated June 28, 2022. Accessed August 1, 2023.

PA DEPARTMENT OF HUMAN SERVICES
MAAC BRIEFING DOCUMENT
DUPIXENT (DUPILUMAB)

4. Wollenberg A, Christen-Zäch S, Taieb A, et al. ETFAD/EADV Eczema task force 2020 position paper on diagnosis and treatment of atopic dermatitis in adults and children. *J Eur Acad Dermatol Venereol*. 2020;34(12):2717-2744.
5. Drucker AM, Ellis AG, Bohdanowicz M, et al. Systemic immunomodulatory treatment for patients with atopic dermatitis – a systemic review and network meta-analysis. *JAMA Dermatol*. 2020;156(6):659-667.
6. Siegels D, Heratizadeh A, Abraham S, et al. Systemic treatments in the management of atopic dermatitis: a systematic review and meta-analysis. *Allergy*. 2021;76(4):1053-1076.
7. Sawangjit R, Dilokthornsakul P, Lloyd-Lavery A, Lai NM, Dellavalle R, Chaiyakunapruk N. Systemic treatments for eczema: a network meta-analysis. *Cochrane Database Syst Rev*. 2020;9:CD013206. Published 2020 Sep 14.
8. Wenzel S. Treatment of severe asthma in adolescents and adults. In: UpToDate [internet database]. Bochner BS, Hollingsworth H, eds. Waltham, MA: UpToDate. Updated March 23, 2022. Accessed April 18, 2022.
9. Global Initiative for Asthma. Global strategy for asthma management and prevention, 2022. Accessed April 18, 2022.
10. Global Initiative for Asthma. Difficult-to-treat & severe asthma in adolescents and adult patients – diagnosis and management, April 2019. Accessed May 3, 2021.
11. U.S. Department of Health, National Institutes of Health, National Heart, Lung, and Blood Institute. Expert panel report 3 (EPR-3): Guidelines for the diagnosis and management of asthma – Full Report 2007. Published October 2007. Accessed April 18, 2022.
12. U.S. Department of Health, National Institutes of Health, National Heart, Lung, and Blood Institute. 2020 Focused Updates to the Asthma Management Guidelines: A Report from the National Asthma Education and Prevention Program Coordinating Committee Expert Panel Working Group. Published December 2020.
13. Hamilos DL, Holbrook EH. Chronic rhinosinusitis: management. In: UpToDate [internet database]. Corren J, Deschler DG, Feldweg AM, eds. Waltham, MA: UpToDate. Updated February 17, 2021. Accessed May 5, 2021.
14. Sidbury R, Alikhan A, Bercovitch L, et al. Guidelines of care for the management of atopic dermatitis in adults with topical therapies. *J Am Acad Dermatol*. 2023;89(1):e1-e20.
15. Yosipovitch G, Mollanazar N, Ständer S, et al. Dupilumab in patients with prurigo nodularis: two randomized, double-blind, placebo-controlled phase 3 trials. *Nat Med*. 2023;29:1180-1190.
16. Ständer S, Pereira MP, Berger T, et al. IFSI-guideline on chronic prurigo including prurigo nodularis. *Itch*. 2020;5(4):e42. doi: 10.1097/itx.0000000000000042.
17. Elmariah S, Kim B, Berger T, et al. Practical approaches for diagnosis and management of prurigo nodularis: United States expert panel consensus. *J Am Acad Dermatol*. 2021;84(3):747-760.
18. **Murrell DF, Ramirez-Quizon M. Management and prognosis of bullous pemphigoid. In: UpToDate [internet database]. Zone JJ, Ofori AO, eds. Waltham, MA: UpToDate. Updated June 23, 2023. Accessed August 26, 2025.**